


**MARKED UP COPY OF AMENDED CLAIMS**  
**(Do not enter)**

1.(TWICE AMENDED)

A compound of the formula



I

or a pharmaceutically acceptable salt thereof, wherein

A is -CR<sub>7</sub>;

B is -NR<sub>1</sub>R<sub>2</sub>, -CR<sub>1</sub>R<sub>2</sub>R<sub>11</sub>, -C(=CR<sub>2</sub>R<sub>12</sub>)R<sub>1</sub>, -NHCHR<sub>1</sub>R<sub>2</sub>, -OCHR<sub>1</sub>R<sub>2</sub>, -SCHR<sub>1</sub>R<sub>2</sub>, -CHR<sub>2</sub>OR<sub>1</sub>, -CHR<sub>1</sub>OR<sub>2</sub>, -CHR<sub>2</sub>SR<sub>1</sub>, -C(S)R<sub>2</sub>, -C(O)R<sub>2</sub>, -CHR<sub>2</sub>NR<sub>1</sub>R<sub>2</sub>, -CHR<sub>1</sub>NHR<sub>2</sub>, -CHR<sub>1</sub>N(CH<sub>3</sub>)R<sub>2</sub>, or -NR<sub>12</sub>NR<sub>1</sub>R<sub>2</sub>;

Z is NH, O, S, -N(C<sub>1</sub>-C<sub>2</sub> alkyl), -NC(O)CF<sub>3</sub>, or -C(R<sub>13</sub>R<sub>14</sub>), wherein R<sub>13</sub> and R<sub>14</sub> are each, independently, hydrogen, trifluoromethyl or methyl, or one of R<sub>13</sub> and R<sub>14</sub> is cyano and the other is hydrogen or methyl, or -C(R<sub>13</sub>R<sub>14</sub>) is a cyclopropyl group, or Z is nitrogen or CH and forms a five or six membered heterocyclic ring fused with R<sub>5</sub>, which ring optionally includes two or three further hetero members selected independently from oxygen, nitrogen, NR<sub>12</sub>, and S(O)<sub>m</sub>, and optionally includes from one to three double bonds, and is optionally substituted with halo, C<sub>1</sub>-C<sub>4</sub> alkyl, -O(C<sub>1</sub>-C<sub>4</sub> alkyl), NH<sub>2</sub>, NHCH<sub>3</sub>, N(CH<sub>3</sub>)<sub>2</sub>, CF<sub>3</sub>, or OCF<sub>3</sub>, with the proviso that said ring does not include any -S-S-, -S-O-, -N-S-, or -O-O- bonds, and does not include more than two oxygen or S(O)<sub>m</sub> heterologous members;

R<sub>1</sub> is C(O)H, C(O)(C<sub>1</sub>-C<sub>6</sub> alkyl), C(O)(C<sub>1</sub>-C<sub>6</sub> alkylene)(C<sub>3</sub>-C<sub>8</sub> cycloalkyl), C(O)(C<sub>3</sub>-C<sub>8</sub> cycloalkylene)(C<sub>3</sub>-C<sub>8</sub> cycloalkyl), C(O)(C<sub>1</sub>-C<sub>6</sub> alkylene)(C<sub>4</sub>-C<sub>8</sub> heterocycloalkyl), -C(O)(C<sub>3</sub>-C<sub>8</sub> cycloalkylene)(C<sub>4</sub>-C<sub>8</sub> heterocycloalkyl), C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>4</sub>-C<sub>8</sub> heterocycloalkyl, -(C<sub>1</sub>-C<sub>6</sub> alkylene)(C<sub>3</sub>-C<sub>8</sub> cycloalkyl), -(C<sub>3</sub>-C<sub>8</sub> cycloalkylene)(C<sub>3</sub>-C<sub>8</sub> cycloalkyl), -(C<sub>1</sub>-C<sub>6</sub> alkylene)(C<sub>4</sub>-C<sub>8</sub> heterocycloalkyl), -(C<sub>3</sub>-C<sub>8</sub> cycloalkylene)(C<sub>4</sub>-C<sub>8</sub> heterocycloalkyl), or -O-aryl, or -O-(C<sub>1</sub>-C<sub>6</sub> alkylene)-aryl; wherein said aryl, C<sub>4</sub>-C<sub>8</sub> heterocycloalkyl, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>3</sub>-C<sub>8</sub> cycloalkylene, and C<sub>1</sub>-C<sub>6</sub> alkylene groups may each independently be optionally substituted with from one to six fluoro and may each independently be optionally substituted with one or two substituents R<sub>8</sub> independently selected from the group consisting of C<sub>1</sub>-C<sub>4</sub> alkyl, -C<sub>3</sub>-C<sub>8</sub> cycloalkyl, hydroxy, chloro, bromo, iodo, CF<sub>3</sub>, -O-(C<sub>1</sub>-C<sub>6</sub> alkyl), -O-(C<sub>3</sub>-C<sub>5</sub> cycloalkyl), -O-CO-(C<sub>1</sub>-C<sub>4</sub> alkyl), -O-CO-NH(C<sub>1</sub>-C<sub>4</sub> alkyl), -O-CO-N(R<sub>24</sub>)(R<sub>25</sub>), -N(R<sub>24</sub>)(R<sub>25</sub>), -S(C<sub>1</sub>-C<sub>4</sub> alkyl), -S(C<sub>3</sub>-C<sub>5</sub> cycloalkyl), -N(C<sub>1</sub>-C<sub>4</sub>alkyl)CO(C<sub>1</sub>-C<sub>4</sub> alkyl), -NHCO(C<sub>1</sub>-C<sub>4</sub> alkyl), -COO(C<sub>1</sub>-C<sub>4</sub> alkyl), -CONH(C<sub>1</sub>-C<sub>4</sub> alkyl), -CON(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), CN, NO<sub>2</sub>, -OSO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl), S<sup>+</sup>(C<sub>1</sub>-C<sub>6</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl)I<sup>-</sup>, -SO(C<sub>1</sub>-C<sub>4</sub> alkyl) and -SO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub>

alkyl); and wherein the C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkylene, C<sub>5</sub>-C<sub>8</sub> cycloalkyl, C<sub>5</sub>-C<sub>8</sub> cycloalkylene, and C<sub>5</sub>-C<sub>8</sub> heterocycloalkyl moieties of R<sub>1</sub> may optionally independently include from one to three double or triple bonds; and wherein the C<sub>1</sub>-C<sub>4</sub> alkyl moieties and C<sub>1</sub>-C<sub>6</sub> alkyl moieties of R<sub>8</sub> can optionally independently be substituted with hydroxy, amino, C<sub>1</sub>-C<sub>4</sub> alkyl, aryl, -CH<sub>2</sub>-aryl, C<sub>3</sub>-C<sub>5</sub> cycloalkyl, or -O-(C<sub>1</sub>-C<sub>4</sub> alkyl), and can optionally independently be substituted with from one to six fluoro, and can optionally include one or two double or triple bonds; and wherein each heterocycloalkyl group of R<sub>1</sub> includes from one to three heteromoieties selected from oxygen, S(O)<sub>m</sub>, nitrogen, and NR<sub>12</sub>;

R<sub>2</sub> is hydrogen, C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>4</sub>-C<sub>8</sub> heterocycloalkyl, -(C<sub>1</sub>-C<sub>6</sub> alkylene)(C<sub>3</sub>-C<sub>8</sub> cycloalkyl), -(C<sub>3</sub>-C<sub>8</sub> cycloalkylene)(C<sub>3</sub>-C<sub>8</sub> cycloalkyl), -(C<sub>1</sub>-C<sub>6</sub> alkylene)(C<sub>4</sub>-C<sub>8</sub> heterocycloalkyl), -(C<sub>3</sub>-C<sub>8</sub> cycloalkylene)(C<sub>4</sub>-C<sub>8</sub> heterocycloalkyl), aryl, -(C<sub>1</sub>-C<sub>6</sub> alkylene)aryl, or -(C<sub>3</sub>-C<sub>8</sub> cycloalkylene)(aryl); wherein each of the foregoing R<sub>2</sub> groups may optionally be substituted with from one to three substituents independently selected from chloro, fluoro, and C<sub>1</sub>-C<sub>6</sub> alkyl, wherein one of said one to three substituents can further be selected from bromo, iodo, C<sub>1</sub>-C<sub>6</sub> alkoxy, -OH, -O-CO-(C<sub>1</sub>-C<sub>6</sub> alkyl), -O-CO-N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -S(C<sub>1</sub>-C<sub>6</sub> alkyl), -S(O)(C<sub>1</sub>-C<sub>6</sub> alkyl), -S(O)<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl), S<sup>+</sup>(C<sub>1</sub>-C<sub>6</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl)<sup>-</sup>, CN, and NO<sub>2</sub>; and wherein the C<sub>1</sub>-C<sub>12</sub> alkyl, -(C<sub>1</sub>-C<sub>6</sub> alkylene), -(C<sub>5</sub>-C<sub>8</sub> cycloalkyl), -(C<sub>5</sub>-C<sub>8</sub> cycloalkylene), and -(C<sub>5</sub>-C<sub>8</sub> heterocycloalkyl) moieties of R<sub>2</sub> may optionally independently include from one to three double or triple bonds; and wherein each heterocycloalkyl group of R<sub>2</sub> includes from one to three heteromoieties selected from oxygen, S(O)<sub>m</sub>, nitrogen, and NR<sub>12</sub>;

or when R<sub>1</sub> and R<sub>2</sub> are as in -NHCHR<sub>1</sub>R<sub>2</sub>, -OCHR<sub>1</sub>R<sub>2</sub>, -SCHR<sub>1</sub>R<sub>2</sub>, -CHR<sub>1</sub>R<sub>2</sub> or -NR<sub>1</sub>R<sub>2</sub>, R<sub>1</sub> and R<sub>2</sub> of B may form a saturated 5- to 8-membered ring which may optionally include one or two double bonds and in which one or two of the ring carbons may optionally be replaced by an oxygen, S(O)<sub>m</sub>, nitrogen or NR<sub>12</sub>; and which ring can optionally be substituted with from 1 to 3 substituents selected from the group consisting of hydroxy, C<sub>1</sub>-C<sub>4</sub> alkyl, fluoro, chloro, bromo, iodo, CF<sub>3</sub>, -O-(C<sub>1</sub>-C<sub>4</sub> alkyl), -O-CO-(C<sub>1</sub>-C<sub>4</sub> alkyl), -O-CO-NH(C<sub>1</sub>-C<sub>4</sub> alkyl), -O-CO-N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -NH(C<sub>1</sub>-C<sub>4</sub> alkyl), -N(C<sub>1</sub>-C<sub>2</sub> alkyl)(C<sub>1</sub>-C<sub>4</sub> alkyl), -S(C<sub>1</sub>-C<sub>4</sub> alkyl), -N(C<sub>1</sub>-C<sub>4</sub> alkyl)CO(C<sub>1</sub>-C<sub>4</sub> alkyl), -NHCO(C<sub>1</sub>-C<sub>4</sub> alkyl), -COO(C<sub>1</sub>-C<sub>4</sub> alkyl), -CONH(C<sub>1</sub>-C<sub>4</sub> alkyl), -CON(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), CN, NO<sub>2</sub>, -OSO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO(C<sub>1</sub>-C<sub>4</sub> alkyl), and -SO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl), wherein one of said one to three substituents can further be selected from phenyl;

R<sub>3</sub> is methyl, ethyl, fluoro, chloro, bromo, iodo, cyano, methoxy, OCF<sub>3</sub>, NH<sub>2</sub>, NH(C<sub>1</sub>-C<sub>2</sub> alkyl), N(CH<sub>3</sub>)<sub>2</sub>, -NHCOCF<sub>3</sub>, -NHCH<sub>2</sub>CF<sub>3</sub>, S(O)<sub>m</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl), CONH<sub>2</sub>, -CONHCH<sub>3</sub>, CON(CH<sub>3</sub>)<sub>2</sub>, -CF<sub>3</sub>, or CH<sub>2</sub>OCH<sub>3</sub>;

R<sub>4</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>3</sub>-C<sub>5</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>4</sub> alkylene)(C<sub>3</sub>-C<sub>5</sub> cycloalkyl), -(C<sub>3</sub>-C<sub>5</sub> cycloalkylene)(C<sub>3</sub>-C<sub>5</sub> cycloalkyl), cyano, fluoro, chloro, bromo, iodo, -OR<sub>24</sub>, C<sub>1</sub>-C<sub>6</sub> alkoxy, -O-(C<sub>3</sub>-C<sub>5</sub> cycloalkyl), -O-(C<sub>1</sub>-C<sub>4</sub> alkylene)(C<sub>3</sub>-C<sub>5</sub> cycloalkyl), -O-(C<sub>3</sub>-C<sub>5</sub> cycloalkylene)(C<sub>3</sub>-C<sub>5</sub> cycloalkyl), -CH<sub>2</sub>SC(S)O(C<sub>1</sub>-C<sub>4</sub> alkyl), -CH<sub>2</sub>OCF<sub>3</sub>, CF<sub>3</sub>, amino, nitro, -NR<sub>24</sub>R<sub>25</sub>, -(C<sub>1</sub>-C<sub>4</sub> alkylene)-OR<sub>24</sub>, -(C<sub>1</sub>-C<sub>4</sub> alkylene)Cl, -(C<sub>1</sub>-C<sub>4</sub> alkylene)NR<sub>24</sub>R<sub>25</sub>, -NHCOR<sub>24</sub>, -NHCONR<sub>24</sub>R<sub>25</sub>, -C=NOR<sub>24</sub>, -NHNOR<sub>24</sub>R<sub>25</sub>, -

$S(O)_mR_{24}$ ,  $-C(O)R_{24}$ ,  $-OC(O)R_{24}$ ,  $-C(O)CN$ ,  $-C(O)NR_{24}R_{25}$ ,  $-C(O)NHNHNR_{24}R_{25}$ , and  $-COOR_{24}$ , wherein the alkyl and alkylene groups of  $R_4$  may optionally independently include one or two double or triple bonds and may optionally independently be substituted with one or two substituents  $R_{10}$  independently selected from hydroxy, amino,  $-NHCOCH_3$ ,  $-NHCOCH_2Cl$ ,  $-NH(C_1-C_2 \text{ alkyl})$ ,  $-N(C_1-C_2 \text{ alkyl})(C_1-C_2 \text{ alkyl})$ ,  $-COO(C_1-C_4 \text{ alkyl})$ ,  $-COOH$ ,  $-CO(C_1-C_4 \text{ alkyl})$ ,  $C_1-C_6$  alkoxy,  $C_1-C_3$  thioalkyl, cyano and nitro, and with one to four substituents independently selected from fluoro and chloro;

$R_5$  is aryl or heteroaryl and is substituted with from one to four substituents  $R_{27}$  independently selected from halo,  $C_1-C_{10}$  alkyl,  $-(C_1-C_4 \text{ alkylene})(C_3-C_8 \text{ cycloalkyl})$ ,  $-(C_1-C_4 \text{ alkylene})(C_4-C_8 \text{ heterocycloalkyl})$ ,  $-(C_3-C_8 \text{ cycloalkyl})$ ,  $-(C_4-C_8 \text{ heterocycloalkyl})$ ,  $-(C_3-C_8 \text{ cycloalkylene})(C_3-C_8 \text{ cycloalkyl})$ ,  $-(C_3-C_8 \text{ cycloalkylene})(C_4-C_8 \text{ heterocycloalkyl})$ ,  $C_1-C_4$  haloalkyl,  $C_1-C_4$  haloalkoxy, nitro, cyano,  $-NR_{24}R_{25}$ ,  $-NR_{24}COR_{25}$ ,  $-NR_{24}CO_2R_{26}$ ,  $-COR_{24}$ ,  $-OR_{25}$ ,  $-CONR_{24}R_{25}$ ,  $-CO(NOR_{22})R_{23}$ ,  $-CO_2R_{26}$ ,  $-C=N(OR_{22})R_{23}$ , and  $-S(O)_mR_{23}$ ; wherein said  $C_1-C_{10}$  alkyl,  $C_3-C_8$  cycloalkyl,  $(C_1-C_4 \text{ alkylene})$ ,  $(C_3-C_8 \text{ cycloalkyl})$ ,  $(C_3-C_8 \text{ cycloalkylene})$ , and  $(C_4-C_8 \text{ heterocycloalkyl})$  groups can be optionally substituted with from one to three substituents independently selected from  $C_1-C_4$  alkyl,  $C_3-C_8$  cycloalkyl,  $(C_1-C_4 \text{ alkylene})(C_3-C_8 \text{ cycloalkyl})$ ,  $-(C_3-C_8 \text{ cycloalkylene})(C_3-C_8 \text{ cycloalkyl})$ ,  $C_1-C_4$  haloalkyl, hydroxy,  $C_1-C_6$  alkoxy, nitro halo, cyano,  $-NR_{24}R_{25}$ ,  $-NR_{24}COR_{25}$ ,  $-NR_{24}CO_2R_{26}$ ,  $-COR_{24}$ ,  $-OR_{25}$ ,  $-CONR_{24}R_{25}$ ,  $CO_2R_{26}$ ,  $-CO(NOR_{22})R_{25}$ , and  $-S(O)_mR_{23}$ ; and wherein two adjacent substituents of the  $R_5$  group can optionally form a 5-7 membered ring, saturated or unsaturated, fused to  $R^5$ , which ring optionally can include one, two, or three heterologous members independently selected from O,  $S(O)_m$ , and N, but not any  $-S-S-$ ,  $-O-O-$ ,  $-S-O-$ , or  $-N-S-$  bonds, and which ring is optionally substituted with  $C_1-C_4$  alkyl,  $C_3-C_8$  cycloalkyl,  $-(C_1-C_4 \text{ alkylene})(C_3-C_8 \text{ cycloalkyl})$ ,  $-(C_3-C_8 \text{ cycloalkylene})(C_3-C_8 \text{ cycloalkyl})$ ,  $C_1-C_4$  haloalkyl, nitro, halo, cyano  $-NR_{24}R_{25}$ ,  $-NR_{24}COR_{25}$ ,  $-NR_{24}CO_2R_{26}$ ,  $-COR_{24}$ ,  $-OR_{25}$ ,  $-CONR_{24}R_{25}$ ,  $CO_2R_{26}$ ,  $-CO(NOR_{22})R_{25}$ , or  $-S(O)_mR_{23}$ ; wherein one of said one to four optional substituents  $R_{27}$  can further be selected from  $-SO_2NH(C_1-C_4 \text{ alkyl})$ ,  $-SO_2NH(C_1-C_4 \text{ alkylene})(C_3-C_8 \text{ cycloalkyl})$ ,  $-SO_2NH(C_3-C_8 \text{ cycloalkyl})$ ,  $-SO_2NH(C_3-C_8 \text{ cycloalkylene})(C_3-C_8 \text{ cycloalkyl})$ ,  $-SO_2N(C_1-C_4 \text{ alkyl})(C_1-C_2 \text{ alkyl})$ ,  $-SO_2NH_2$ ,  $-NHSO_2(C_1-C_4 \text{ alkyl})$ ,  $-NHSO_2(C_3-C_8 \text{ cycloalkyl})$ ,  $-NHSO_2(C_1-C_4 \text{ alkylene})(C_3-C_8 \text{ cycloalkyl})$ , and  $-NHSO_2(C_3-C_8 \text{ cycloalkylene})(C_3-C_8 \text{ cycloalkyl})$ ; and wherein the alkyl, and alkylene groups of  $R_5$  may independently optionally include one double or triple bond;

$R_6$  is hydrogen,  $C_1-C_6$  alkyl,  $C_3-C_8$  cycloalkyl,  $-(C_1-C_6 \text{ alkylene})(C_3-C_8 \text{ cycloalkyl})$ , or  $-(C_3-C_8 \text{ cycloalkylene})(C_3-C_8 \text{ cycloalkyl})$ , wherein said alkyl and cycloalkyl may optionally be substituted with one hydroxy, methoxy, ethoxy or fluoro group;

$R_7$  is hydrogen, methyl, fluoro, chloro, bromo, iodo, cyano, hydroxy,  $-O(C_1-C_2 \text{ alkyl})$ ,  $-O(\text{cyclopropyl})$ ,  $-COO(C_1-C_2 \text{ alkyl})$ ,  $-COO(C_3-C_8 \text{ cycloalkyl})$ ,  $-OCF_3$ ,  $CF_3$ ,  $-CH_2OH$ , or  $CH_2OCH_3$ ;

$R_{11}$  is hydrogen, hydroxy, fluoro, ethoxy, or methoxy;

$R_{12}$  is hydrogen or  $C_1-C_4$  alkyl;

$R_{22}$  is independently at each occurrence selected from hydrogen,  $C_1-C_4$  alkyl,  $C_1-C_4$

haloalkyl, C<sub>3</sub>-C<sub>6</sub> alkenyl, C<sub>3</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, (C<sub>3</sub>-C<sub>8</sub> cycloalkylene)(C<sub>3</sub>-C<sub>8</sub> cycloalkyl), and (C<sub>1</sub>-C<sub>4</sub> alkylene)(C<sub>3</sub>-C<sub>8</sub> cycloalkyl);

R<sub>23</sub> is independently at each occurrence selected from C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, C<sub>2</sub>-C<sub>8</sub> alkoxyalkyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>4</sub> alkylene)(C<sub>3</sub>-C<sub>8</sub> cycloalkyl), -(C<sub>3</sub>-C<sub>8</sub> cycloalkylene)(C<sub>3</sub>-C<sub>8</sub> cycloalkyl), aryl, -(C<sub>1</sub>-C<sub>4</sub> alkylene)aryl, piperidine, pyrrolidine, piperazine, N-methylpiperazine, morpholine, and thiomorpholine;

R<sub>24</sub> and R<sub>25</sub> are independently at each occurrence selected from hydrogen, -C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, especially CF<sub>3</sub>, -CHF<sub>2</sub>, CF<sub>2</sub>CF<sub>3</sub>, or CH<sub>2</sub>CF<sub>3</sub>, -(C<sub>1</sub>-C<sub>4</sub> alkylene)OH, -(C<sub>1</sub>-C<sub>4</sub> alkylene)-O-(C<sub>1</sub>-C<sub>4</sub> alkyl), -(C<sub>1</sub>-C<sub>4</sub> alkylene)-O-(C<sub>3</sub>-C<sub>5</sub> cycloalkyl), C<sub>3</sub>-C<sub>8</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>4</sub> alkylene)(C<sub>3</sub>-C<sub>8</sub> cycloalkyl), -(C<sub>3</sub>-C<sub>8</sub> cycloalkylene)(C<sub>3</sub>-C<sub>8</sub> cycloalkyl), -C<sub>4</sub>-C<sub>8</sub> heterocycloalkyl, -(C<sub>1</sub>-C<sub>4</sub> alkylene)(C<sub>4</sub>-C<sub>8</sub> heterocycloalkyl), -(C<sub>3</sub>-C<sub>8</sub> cycloalkylene)(C<sub>4</sub>-C<sub>8</sub> heterocycloalkyl), aryl, and -(C<sub>1</sub>-C<sub>4</sub> alkylene)(aryl), wherein the -C<sub>4</sub>-C<sub>8</sub> heterocycloalkyl groups can each independently optionally be substituted with aryl, CH<sub>2</sub>-aryl, or C<sub>1</sub>-C<sub>4</sub> alkyl, and can optionally include one or two double or triple bonds; or, when R<sub>24</sub> and R<sub>25</sub> are as NR<sub>24</sub>R<sub>25</sub>, -C(O)NR<sub>24</sub>R<sub>25</sub>, -(C<sub>1</sub>-C<sub>4</sub> alkylene)NR<sub>24</sub>R<sub>25</sub>, or -NHCONR<sub>24</sub>R<sub>25</sub>, then NR<sub>24</sub>R<sub>25</sub> may further optionally form a 4 to 8 membered heterocyclic ring optionally including one or two further hetero members independently selected from S(O)<sub>m</sub>, oxygen, nitrogen, and NR<sub>12</sub>, and optionally including from one to three double bonds;

R<sub>26</sub> is independently at each occurrence selected from C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> haloalkyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, -(C<sub>1</sub>-C<sub>4</sub> alkylene)(C<sub>3</sub>-C<sub>8</sub> cycloalkyl), -(C<sub>3</sub>-C<sub>8</sub> cycloalkylene)(C<sub>3</sub>-C<sub>8</sub> cycloalkyl), aryl, and -(C<sub>1</sub>-C<sub>4</sub> alkylene)(aryl); and

wherein each m is independently zero, one, or two,

with the proviso that heterocycloalkyl groups of the compound of formula I~~[I, II, or III]~~ do not include any -S-S-, -S-O-, -N-S-, or -O-O- bonds, and do not include more than two oxygen or S(O)<sub>m</sub> heterologous members.

9. (AMENDED) A pharmaceutical composition for the treatment of (a) a disorder or condition the treatment of which can be effected or facilitated by antagonizing CRF, or (b) a disorder or condition selected from inflammatory disorders such as rheumatoid arthritis and osteoarthritis, pain, asthma, psoriasis and allergies; generalized anxiety disorder; panic; phobias, including social phobia, agoraphobia, and specific phobias; obsessive-compulsive disorder; post-traumatic stress disorder; sleep disorders induced by stress; pain perception such as fibromyalgia; mood disorders such as depression, including major depression, single episode depression, recurrent depression, child abuse induced depression, mood disorders associated with premenstrual syndrome, and postpartum depression; dysthemia; bipolar disorders; cyclothymia; chronic fatigue syndrome; stress-induced headache; ~~{cancer;}~~ irritable bowel syndrome~~[- Crohn's disease];~~ spastic colon; post operative ileus; ulcer; diarrhea; stress-induced fever; ~~[human immunodeficiency virus infections;]~~ neurodegenerative diseases such as

Alzheimer's disease, Parkinson's disease and Huntington's disease; gastrointestinal diseases; ~~[eating disorders such as anorexia and bulimia nervosa;]~~ hemorrhagic stress; chemical dependencies or addictions, including dependencies or addictions to alcohol, cocaine, heroin, benzodiazapines, or other drugs; drug or alcohol withdrawal symptoms; stress-induced psychotic episodes; euthyroid sick syndrome; syndrome of inappropriate antidiuretic hormone; ~~[obesity; infertility;]~~ head trauma; spinal cord trauma; ischemic neuronal damage, including cerebral ischemia, for example cerebral hippocampal ischemia; excitotoxic neuronal damage; epilepsy; stroke; immune dysfunctions including stress induced immune dysfunctions, including porcine stress syndrome, bovine shipping fever, equine paroxysmal fibrillation, confinement dysfunction in chicken, sheering stress in sheep, and human-animal interaction stress in dogs; muscular spasms; urinary incontinence; senile dementia of the Alzheimer's type; multiinfarct dementia; amyotrophic lateral sclerosis; hypertension; tachycardia; congestive heart failure; osteoporosis~~[-]~~ and premature birth~~[- hypoglycemia, and Syndrome X]~~ in a mammal or bird, comprising an amount of a compound according to claim 1 that is effective in the treatment of such disorder or condition, and a pharmaceutically acceptable carrier.

10. (AMENDED) A method for the treatment of (a) a disorder or condition the treatment of which can be effected or facilitated by antagonizing CRF, or (b) a disorder or condition selected from inflammatory disorders such as rheumatoid arthritis and osteoarthritis, pain, asthma, psoriasis and allergies; generalized anxiety disorder; panic; phobias, including social phobia, agoraphobia, and specific phobias; obsessive-compulsive disorder; post-traumatic stress disorder; sleep disorders induced by stress; pain perception such as fibromyalgia; mood disorders such as depression, including major depression, single episode depression, recurrent depression, child abuse induced depression, mood disorders associated with premenstrual syndrome, and postpartum depression; dysthemia; bipolar disorders; cyclothymia; chronic fatigue syndrome; stress-induced headache; ~~[cancer;]~~ irritable bowel syndrome~~[- Crohn's disease]~~; spastic colon; post operative ileus; ulcer; diarrhea; stress-induced fever; ~~[human immunodeficiency virus infections;]~~ neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease and Huntington's disease; gastrointestinal diseases; ~~[eating disorders such as anorexia and bulimia nervosa;]~~ hemorrhagic stress; chemical dependencies or addictions, including dependencies or addictions to alcohol, cocaine, heroin, benzodiazapines, or other drugs; drug or alcohol withdrawal symptoms; stress-induced psychotic episodes; euthyroid sick syndrome; syndrome of inappropriate antidiuretic hormone; ~~[obesity; infertility;]~~ head trauma; spinal cord trauma; ischemic neuronal damage, including cerebral ischemia, for example cerebral hippocampal ischemia; excitotoxic neuronal damage; epilepsy; stroke; immune dysfunctions including stress induced immune dysfunctions, including porcine stress syndrome, bovine shipping fever, equine paroxysmal fibrillation, confinement dysfunction in chicken, sheering stress in sheep, and human-animal interaction stress in dogs; muscular spasms; urinary incontinence;

senile dementia of the Alzheimer's type; multiinfarct dementia; amyotrophic lateral sclerosis; hypertension; tachycardia; congestive heart failure; osteoporosis[;] and premature birth[; hypoglycemia, and Syndrome X] in a mammal or bird, comprising administering to a subject in need of said treatment an amount of a compound according to claim 1, that is effective in treating such disorder or condition.

17. A pharmaceutical composition for treating a condition comprising a compound of claim 1 in an amount effective to treat said condition and a pharmaceutically acceptable carrier, wherein said condition is selected from the group consisting of:

- a) abnormal circadian rhythm;
- b) depression, further wherein a second compound for treating depression is administered, said second compound for treating depression having an onset of action that is delayed with respect to that of said ~~[CRF-antagonist]~~ compound of claim 1; and
- c) emesis.